


Poison - Chocolate



1 Try to identify the toxic substance collect any empty bottles or containers, and check for distinctive odors



2 Look for traces in or around the mouth to identify the substance



3 If your dog vomits, note carefully the appearance of the vomit and collect them in a plastic baggy for identification



4 Try to estimate the amount that the pet has eaten. Some things are only toxic in large doses



5 If your dog ingested the chocolate less than 2 hours ago induce vomiting with 3% Hydrogen Peroxide, 1-2 teaspoons for every 10lb your dog weighs, and squirt it at the back of your dog's tongue



6 You will notice a foaming action and the taste should prompt vomiting within 5 min, if needed repeat 2-3 times allowing 5 min between doses



7 Call the ASPCA Animal Poison Control Center hotline which operates 24 hours a day, 7 days a week, at (888) 426-4435 there is a \$60 consultation fee, payable by credit card, for this service and this includes follow-up consultation should you or your vet need further assistance with your pet's case

- ⚠ Induce vomiting with 3% Hydrogen Peroxide
- ⚠ Wear gloves while helping your dog to avoid contact with the acid
- ⚠ A dose of less than 1 oz of milk chocolate per pound of body weight could potentially cause death
- ⚠ If your animal is having seizures, losing consciousness, is unconscious or is having difficulty breathing, telephone ahead and bring your pet immediately to your local veterinarian or emergency veterinary clinic
- ✅ Note any symptoms that your pet may be having
- ✅ ASPCA Animal Poison Control Center hotline which operates 24 hours a day, 7 days a week, at (888) 426-4435
- ✅ You can give activated charcoal by mouth to help absorb antifreeze that was recently ingested but call your veterinarian or the ASPCA Animal Poison Control Center hotline first.