


Poison - Pain Medicine Aspirin, Acetaminophen



1 Try to identify the toxic substance collect any empty bottles or containers, and check for distinctive odors



2 Look for traces in or around the mouth to identify the substance



3 If your dog vomits, note carefully the appearance of the vomit and collect them in a plastic baggy for identification



4 Try to estimate the amount that the pet has eaten. Some things are only toxic in large doses



5 If your dog ingested the poison less than 2 hours ago induce vomiting with 3% Hydrogen Peroxide, 1-2 teaspoons for every 10lb your dog weighs, and squirt it at the back of your dog's tongue



6 You will notice a foaming action and the taste should prompt vomiting within 5 min, if needed repeat 2-3 times allowing 5 min between doses



7 Call the ASPCA Animal Poison Control Center hotline which operates 24 hours a day, 7 days a week, at (888) 426-4435 there is a \$60 consultation fee, payable by credit card, for this service and this includes follow-up consultation should you or your vet need further assistance with your pet's case



Induce vomiting with 3% Hydrogen Peroxide



If your animal is having seizures, losing consciousness, is unconscious or is having difficulty breathing, telephone ahead and bring your pet immediately to your local veterinarian or emergency veterinary clinic



Note any symptoms that your pet may be having



ASPCA Animal Poison Control Center hotline which operates 24 hours a day, 7 days a week, at (888) 426-4435



You can give activated charcoal by mouth to help absorb chocolate that was recently ingested but call your veterinarian or the ASPCA Animal Poison Control Center hotline first.